



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

By substituting the values of x found above in the original equation, we see that 0 and $\pm \infty$ are to be rejected. We find, however, by inspection that $x = 1$ is a root.

Hence, the roots are 1 and $(1 \pm \sqrt{5})/2$.

Also solved by ALBERT N. NAUER, A. M. HARDING, C. E. GITHENS, V. M. SPUNAR, ELIJAH SWIFT, W. C. EELLS, G. W. HARTWELL, and the PROPOSER.

430A. Proposed by H. C. FEEMSTER, York College, Neb.

Solve the equations

$$\sum_{i=1}^n x_i - x_n = k + \frac{n^2 - 3n + 2}{2} d, \quad (1)$$

$$\sum_{i=1}^n x_i - x_{n-1} = k + \frac{n^2 - 3n + 4}{2} d, \quad (2)$$

$$\sum_{i=1}^n x_i - x_{n-2} = k + \frac{n^2 - 3n + 6}{2} d, \quad (3)$$

$$\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \vdots$$

$$\sum_{i=1}^n x_i - x_1 = k + \frac{n^2 - n}{2} d. \quad (n)$$

SOLUTION BY A. M. HARDING, Univ. of Arkansas.

Add the given equations and obtain

$$(n-1) \sum_{i=1}^n x_i = nk + \frac{n^3 - 3n^2 + n^2 + n}{2} d,$$

or

$$\sum_{i=1}^n x_i = \frac{n}{n-1} \cdot k + \frac{n(n-1)}{2} d.$$

Subtract each of the given equations from this equation and obtain

$$x_n = \frac{k}{n-1} + (n-1)d,$$

$$x_{n-1} = \frac{k}{n-1} + (n-2)d,$$

$$x_{n-2} = \frac{k}{n-1} + (n-3)d,$$

$$\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot$$

$$x_2 = \frac{k}{n-1} + d,$$

$$x_1 = \frac{k}{n-1}.$$

Also solved by NATHAN ALTSHILLER, S. A. JOFFE, J. W. CLAWSON, FRANK R. MORRIS, ELBERT H. CLARKE, HORACE OLSON, N. P. PANDYA, and the PROPOSER.

431. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.

Form a magic square of 9 cells such that (the integers being all different) the products of the integers in the rows, columns, and diagonals shall be the same and the smallest product possible.

SOLUTION BY S. A. JOFFE, New York City.

All the nine integers will be determined if, in addition to the product p of three integers of any row (column, or diagonal), we know two extremes, say of the first row, a and b , and the central integer c .

| | | |
|-----|-----|-----|
| a | | b |
| | c | |
| | | |

The extremes of the third row will then be p/ac and p/bc ; of the second row will be bc/a and ac/b ; and those of the second column will be p/ab and ab/c .

Hence the product of the integers in the second row equals

$$\frac{bc}{a} \cdot c \cdot \frac{ac}{b} = c^3,$$

and therefore

$$p = c^3,$$

so that the magic square becomes

| | | |
|-----------------|------------------|-----------------|
| a | $\frac{c^3}{ab}$ | b |
| $\frac{bc}{a}$ | c | $\frac{ac}{b}$ |
| $\frac{c^3}{b}$ | $\frac{ab}{c}$ | $\frac{c^3}{a}$ |

If we disregard the central integer c , the remaining eight integers must therefore form four pairs: (1) m_1 and c^2/m_1 ; (2) m_2 and c^2/m_2 ; (3) m_3 and c^2/m_3 ; (4) m_4 and c^2/m_4 . Taking now for m_1, m_2, m_3 and m_4 the smallest four integers, i. e., 1, 2, 3 and 4, we find that the *smallest square* number c^2 , divisible by these four integers, is 36, and consequently $c = \sqrt{36} = 6$.

However, no corner number, say a , can be unity; because, if $a = 1$ then ac/b and ab/c become c/b and b/c , which must be simultaneously integers, and this is impossible unless $b = c$; but the latter equality is excluded by the condition of the problem. Taking $a = 2$, $b = 3$ and $c = 6$, we find the required magic square with the smallest product possible to be

| | | |
|----|----|----|
| 2 | 36 | 3 |
| 9 | 6 | 4 |
| 12 | 1 | 18 |

Also solved by HERBERT N. CARLETON, N. P. PANDYA.